

Hyperon semileptonic decay constants and quark-spin content of the octet baryons

Hyun-Chul Kim^a, Michał Przaszłowicz^b and Klaus Goeke^c

^aDepartment of Physics, Pusan National University,
609-735 Pusan, Republic of Korea

^bInstitute of Physics, Jagellonian University,
Cracow, Poland

^cInstitute for Theoretical Physics II, Ruhr-University Bochum,
D-44780 Bochum, Germany

We present the recent investigation on the effect of the SU(3) symmetry breaking in the spin structure of the octet baryons.

The spin structure of the nucleon has been a hot issue since the measurement of the first moment of the proton spin structure function g_1^p [1] by the European Muon Collaboration (EMC). The EMC result indicates that the strange quark is polarized in the opposite direction to other light quarks. It implies that the spin of the nucleon is not carried by quarks only. While it is now known that a large fraction of the nucleon spin is provided by gluons and their orbital angular momenta, it is still very important to understand the mechanism of how the quarks carry the nucleon spin. The information on the nucleon spin carried by quarks comes from two different sources: the first moment of the spin structure function $g_1^p(x)$ of the proton and the experimental data of hyperon semileptonic decay (HSD). With SU(3) symmetry assumed, one obtains two reduced matrix elements F and D . Taking for $F = 0.46$ and for $D = 0.80$, one gets for the proton: $\Delta u_p = 0.79$, $\Delta d_p = -0.47$ and $\Delta s_p = -0.13$, which implies $\Delta \Sigma_p = 0.19$, quite a small number as compared with the naive expectation from the quark model: $\Delta \Sigma_p = 1$.

Recently, the spin content of the Λ hyperon has drawn attention to us. In contrast to the nucleon, the nonrelativistic quark model predicts that the spin of the Λ comes solely from the strange quark, since the up and down quarks constitute the spin singlet. Furthermore, the Δq_Λ is deeply related to the fragmentation function which can be measured experimentally [2].

However, note that $\Delta \Sigma_p$ and $\Delta \Sigma_\Lambda$ are *not directly measured*. The standard way to extract $\Delta \Sigma_B$ is to assume the SU(3) symmetry for HSD. In this case it is enough to take *any* two decays and data of deep inelastic scattering as input. With SU(3) symmetry assumed, one can pick out two different values of the constants of HSD among the existing data. One normally uses neutron and $\Sigma^- \rightarrow \beta$ decays as an input. However, if the SU(3)

symmetry breaking were not important, any pair out of the six known HSD constants would give roughly the same number for $\Delta\Sigma_B$. This is, however, *not* the case. In fact, the effect of SU(3) symmetry breaking in HSD is not at all small. Furthermore, the uncertainty of the HSD constant for $\Xi^- \rightarrow \Sigma^0$ is noticeably larger than those for the other processes. Thus, it is still of great significance to understand the effect of SU(3) symmetry breaking on the quark-spin content of the nucleon.

In order to analyze the symmetry breaking in weak decays, it is inevitable to resort to some specific model [3]. We employ the chiral quark-soliton model(χ QSM): See Ref.[4] and references therein) to incorporate the SU(3) symmetry breaking into the semileptonic decays. The χ QSM satisfactorily describes the axial-vector properties of hyperons (see Ref. [5] and references therein.). Since the symmetry breaking pattern of the χ QSM is identical to the one derived in large N_c QCD [6], our analysis is in fact much more general than the model itself. In addition, we employ the experimental data for HSD in order to fix the dynamical variables in the model, so that we can minimize the model-dependence of the present study.

However, since $g_A^{(0)}(B)$ does not correspond to the SU(3) octet axial-vector current, a further assumption is required to connect it to the octet axial-vector constants. The χ QSM (as most of the hedgehog models) has a remarkable virtue of connecting the $g_A^{(0)}(B)$ with $g_A^{(3)}$ and $g_A^{(8)}$, and the semileptonic decay constants in a direct manner. This connection introduces a model dependence into our analysis. However, there is no significant numerical difference between the results obtained with and without this model dependent ingredient [7]. Thus, we would like to report in brief the recent investigation on the effect of the SU(3) symmetry breaking, based on Refs. [7,8], to which we refer for details.

Figure 1 shows the dependence of the integrated quark densities Δq_n of the nucleon on the strange quark mass m_s , of which definitions can be found in Ref. [7]:




Figure 1: The dependence of the Δq_n on the strange quark mass m_s .

While the up and down quark densities have smaller errors as m_s increases, the uncertainty of the Δs_n becomes much larger. With $m_s = 180$ MeV, we obtain $\Delta s_n = 0.33 \pm 0.51$. It indicates that Δs_n is in practice undetermined. The reason is that the error of the semileptonic decay constant for the process $\Xi^- \rightarrow \Sigma^0$ propagates strongly in the Δs_n .

Figure 2 plots the flavor decomposition of the quark-spin content of the Λ . As in the case of the nucleon, the error of the Δs_Λ is noticeable with m_s increased.




Figure 2: The dependence of the Δq_Λ on the strange quark mass m_s .

Figure 3 depicts the splitting of the $\Delta\Sigma_B$ s due to the strange quark mass. The result is according to the following equation:

$$g_B^{(0)} \hat{S}_3 = a_3 \hat{S}_3 + \sqrt{3} m_s (a_5 - a_6) \langle B | D_{83}^{(8)} | B \rangle, \quad (1)$$

where $\hat{S}_3$ is the third component of the spin operator. The m_s stands for the strange quark mass which plays a role of breaking SU(3) symmetry. $\langle B | D_{83}^{(8)} | B \rangle$ is the matrix element of the SU(3) Wigner function. The a_i denote the dynamical variables from the model which are fixed by the HSD data. They appear also in the octet axial-vector constants, which makes it possible to link the octet axial-vector constants to the octet one.




Figure 3: Effect of SU(3) symmetry breaking on the $\Delta\Sigma_B$.

Actually, the simplest version of the Sykrme model without vector mesons gives $a_3 = 0$. In the case of the χ QSM provides nonzero a_3 . In the limit of the large soliton, the ratio of a_3/a_i goes to zero, which is in agreement with Ref. [9]. When the size of the soliton becomes closer to zero, which corresponds to the limit of the nonrelativistic quark model,

$g_p^{(0)}$ turns out to be one, as it should be. The removal of the degeneracy in $\Delta\Sigma_{BS}$ comes from the second term in Eq.(1). While the $\Delta\Sigma_p$ and $\Delta\Sigma_\Lambda$ show relatively mild dependence on m_s , the $\Delta\Sigma_\Xi$ and $\Delta\Sigma_\Sigma$ depend quite strongly on m_s . Large error bars with m_s is due to the propagation of the uncertainty in Ξ decay to the variables a_i . From figure 3, we find a very interesting thing. With the effect of the SU(3) symmetry breaking, $\Delta\Sigma_\Lambda$ and $\Delta\Sigma_\Xi$ get larger and finally quite closer to one. It indicates that the spins of Λ and Ξ are almost carried by quarks. That is, Λ and Ξ behave more “nonrelativistically”. Hence, extracting the $\Delta\Sigma_\Lambda$ from the experimental data as precisely as possible and comparing it with that of the nucleon, one would see the effect of the SU(3) symmetry breaking, explicitly.

In summary, we have presented in this talk a recent investigation on the effect of the SU(3) symmetry breaking within the framework of the chiral quark-soliton model, incorporating the experimental data of hyperon semileptonic decay. Because of the large uncertainty in the data for the Ξ decay, one is not able to determine the $\Delta s_{n,\Lambda}$, which means that one can not tell for sure about how much the strange quark carries the nucleon spin.

The SU(3) symmetry breaking removes the degeneracy of the $\Delta\Sigma_{BS}$. Therefore, knowing the difference between the $\Delta\Sigma_p$ and $\Delta\Sigma_\Lambda$ will bring out the effect of the SU(3) symmetry breaking.

The Λ and Ξ behave more “nonrelativistically” than the nucleon.

In order to determine the spin content of the SU(3) baryons carried by quarks, it is highly required to reduce the experimental data of Ξ semileptonic decay.

Acknowledgments

The work of H.C.K has been supported by the Research Fund of Pusan National University. The work of M.P. has been supported by Polish KBN Grant PB 2 P03B 019 17. The work of K.G. has been supported by the BMBF, the DFG, and the COSY–Project(Jülich).

REFERENCES

1. EMC Collaboration, J. Ashman *et al.*, Phys. Lett. **B206** (1988) 364; Nucl.Phys. **B328** (1989) 1.
2. M. Burkardt and R.L. Jaffe, Phys. Rev. Lett. **70** (1993) 2537.
3. M. Karliner and H. Lipkin, Phys. Lett. **B461** (1999)280.
4. C.V. Christov, A. Blotz, H.-Ch. Kim, P. Pobylitsa, T. Watabe, Th. Meissner, E. Ruiz Arriola, and K. Goeke, Prog. Part. Nucl. Phys. **37** (1996) 91.
5. H.-Ch. Kim, M.V. Polyakov, M. Praszalowicz, and K. Goeke, Phys. Rev. **D57** (1998), 299.
6. J. Dai, R. Dasher, E. Jenkins, and A. V. Manohar, Phys. Rev. **D53** (1996) 273; R. Flores-Mendieta, E. Jenkins, and A. V. Manohar, Phys. Rev. **D58** (1998) 94028.
7. H.-Ch. Kim, M. Praszalowicz, and K. Goeke, Phys. Rev. **D61** (2000) 114006.
8. H.-Ch. Kim, M. Praszalowicz and K. Goeke, Acta Phys. Polon. **B31** (2000) 1767; H.-Ch. Kim, M. Praszalowicz and K. Goeke, hep-ph/0007022.
9. S.J. Brodsky, J. Ellis, and M. Karliner, Phys. Lett. **206B** (1988) 309.